

Anästhesie Journal 1/2023

Journal d'anesthésie 1/2023

Referenzen

Références

Kabelloses Monitoring bei postoperativen Patient:innen: Präventiv handeln – IPS-Aufnahmen verringern

Surveillance sans fil des patients postopératoires: Agir de manière préventive – Réduire les admissions aux soins intensifs

Benjamin Albiez

1. Sessler DI, Saugel B. Beyond ‘failure to rescue’: the time has come for continuous ward monitoring. *Br J Anaesth* 2019; 122: 304e6
2. Subbe CP. Validation of a modified early warning score in medical admissions. *QJM* 2001; 94: 521e6
3. McGloin H, Adam SK, Singer M. Unexpected deaths and referrals to intensive care of patients on general wards. Are some cases potentially avoidable? *J R Coll Physicians Lond* 1999; 33
4. Morgan R, Williams F, Wright M. An early warning scoring system for detecting developing critical illness. *Clin Intensive Care* 1997; 8: 100
5. Prytherch DR, Smith GB, Schmidt PE, Featherstone PI. ViEWS – towards a national early warning score for detecting adult inpatient deterioration. *Resuscitation* 2010; 81: 932e7
6. Stenhouse C, Coates S, Tivey M, Allsop P, Parker T. Prospective evaluation of a modified Early Warning Score to aid earlier detection of patients developing critical illness on a general surgical ward. *Br J Anaesth* 2000; 84: 663P
7. Smith MEB, Chiovaro JC, O’Neil M, et al. Early warning system scores for clinical deterioration in hospitalized patients: a systematic review. *Ann Am Thorac Soc* 2014; 11: 1454e65
8. Ludikhuize J, Brunsvelde-Reinders AH, Dijkgraaf MGW, et al. Outcomes associated with the nationwide introduction of rapid response systems in The Netherlands. *Crit Care Med* 2015; 43: 2544e51
9. Alam N, Hobbelink EL, van Tienhoven AJ, van de Ven PM, Jansma EP, Nanayakkara PWB. The impact of the use of the Early Warning Score (EWS) on patient outcomes: a systematic review. *Resuscitation* 2014; 85: 587e94
10. Jones S, Mullally M, Ingleby S, Buist M, Bailey M, Eddleston JM. Bedside electronic capture of clinical observations and automated clinical alerts to improve compliance with an Early Warning Score protocol. *Crit Care Resusc* 2011; 13: 83e8
11. Smith AF, Oakey RJ. Incidence and significance of errors in a patient “track and trigger” system during an epidemic of Legionnaires’ disease: retrospective casenote analysis. *Anaesthesia* 2006; 61: 222e8
12. Hands C, Reid E, Meredith P, et al. Patterns in the recording of vital signs and early warning scores: compliance with a clinical escalation protocol. *BMJ Qual Saf* 2013; 22: 719e26
13. Eddahchouri Y, Koeneman M, Plokker M, et al. Low compliance to a vital sign safety protocol on general hospital wards: a retrospective cohort study. *Int J Nurs Stud* 2021; 115: 103849
14. Michard F, Sessler DI. Ward monitoring 3.0. *Br J Anaesth* 2018; 121: 999e1001
15. Michard F, Kalkman CJ. Rethinking patient surveillance on hospital wards. *Anesthesiology* 2021; 135: 531e40
16. Weenk M, Bredie SJ, Koeneman M, Hesselink G, van Goor H, van de Belt TH. Continuous monitoring of vital signs at the general ward using wearable devices: a qualitative analysis. *J Med Internet Res* 2020; 22, e15471

17. Downey CL, Chapman S, Randell R, Brown JM, Jayne DG. The impact of continuous versus intermittent vital signs monitoring in hospitals: a systematic review and narrative synthesis. *Int J Nurs Stud* 2018; 84: 19e27
18. Sun L, Joshi M, Khan SN, Ashrafi H, Darzi A. Clinical impact of multi-parameter continuous non-invasive monitoring in hospital wards: a systematic review and meta-analysis. *J R Soc Med* 2020; 113: 217e24
19. Weenk M, van Goor H, Frietman B, et al. Continuous monitoring of vital signs using wearable devices on the general ward: pilot study. *JMIR mHealth uHealth* 2017; 5: e91
20. Prgomet M, Cardona-Morrell M, Nicholson M, et al. Vital signs monitoring on general wards: clinical staff perceptions of current practices and the planned introduction of continuous monitoring technology. *Int J Qual Health Care* 2016; 28: 515e21
21. Weenk M, Koeneman M, van de Belt TH, Engelen LJPG, van Goor H, Bredie SJH. Wireless and continuous monitoring of vital signs in patients at the general ward. *Resuscitation* 2019; 136: 47e53
22. Turan A, Chang C, Cohen B, et al. Incidence, severity, and detection of blood pressure perturbations after abdominal surgery. *Anesthesiology* 2019; 130: 550e9
23. Saab R, Wu BP, Rivas E, et al. Failure to detect ward hypoxaemia and hypotension: contributions of insufficient assessment frequency and patient arousal during nursing assessments. *Br J Anaesth* 2021; 127: 760e8
24. Weller RS, Foard KL, Harwood TN. Evaluation of a wireless, portable, wearable multi-parameter vital signs monitor in hospitalized neurological and neurosurgical patients. *J Clin Monit Comput* 2018; 32: 945e51
25. Verrillo SC, Cvach M, Hudson KW, Winters BD. Using continuous vital sign monitoring to detect early deterioration in adult postoperative inpatients. *J Nurs Care Qual* 2019; 34: 107e13
26. Taenzer AH, Pyke JB, McGrath SP, Blike GT. Impact of pulse oximetry surveillance on rescue events and intensive care unit transfers. *Anesthesiology* 2010; 112: 282e7
27. Brown H, Terrence J, Vasquez P, Bates DW, Zimlichman E. Continuous monitoring in an inpatient medical-surgical unit: a controlled clinical trial. *Am J Med* 2014; 127: 226e32
28. Watkinson PJ, Barber VS, Price JD, Hann A, Tarassenko L, Young JD. A randomised controlled trial of the effect of continuous electronic physiological monitoring on the adverse event rate in high risk medical and surgical patients. *Anaesthesia* 2006; 61: 1031e9
29. Fieselmann JF, Hendryx MS, Helms CM, Wakefield DS. Respiratory rate predicts cardiopulmonary arrest for internal medicine inpatients. *J Gen Intern Med* 1993; 8: 354e60
30. Cardona-Morrell M, Prgomet M, Turner RM, Nicholson M, Hillman K. Effectiveness of continuous or intermittent vital signs monitoring in preventing adverse events on general wards: a systematic review and meta-analysis. *Int J Clin Pract* 2016; 70: 806e24
31. Subbe CP, Duller B, Bellomo R. Effect of an automated notification system for deteriorating ward patients on clinical outcomes. *Crit Care* 2017; 21: 52
32. Boer C, Touw HR, Loer SA. Postanesthesia care by remote monitoring of vital signs in surgical wards. *Curr Opin Anaesthesiology* 2018; 31: 716e22
33. Flick M, Saugel B. Continuous ward monitoring: the selection, monitoring, alarms, response, treatment (SMART) road map. *Br J Anaesth* 2021; 127: 675e7
34. Nedel WL, Silveira F da. Different research designs and their characteristics in intensive care. *Rev Bras Ter Intensiva* 2016; 28: 256e60
35. Kang MA, Churpek MM, Zadravec FJ, Adhikari R, Twu NM, Edelson DP. Real-time risk prediction on the wards. *Crit Care Med* 2016; 44: 1468e73
36. Joanna Briggs Institute, J. (2017). © Joanna Briggs Institute 201. Critical Appraisal Checklist for Analytical Cross Sectional Studies. https://jbi.global/sites/default/files/2019-05/JBI_Critical_Appraisal-Checklist_for_Analytical_Cross_Sectional_Studies2017_0.pdf